

ENTEROTOXAEMIA VACCINE IP

ET VAC[®]

ईटी वैक

COMPOSITION

Each dose of ET VAC[®], Enterotoxaemia Vaccine contains formalin killed cultures of *Clostridium perfringens* type D and Epsilon toxoid, adjuvanted with aluminium hydroxide gel, having potency of not less than 2 IU of epsilon antitoxin per mL of pooled serum. Formaldehyde ($\leq 0.05\%$ w/v) is added as a preservative.

INDICATIONS

ET VAC[®] is used for prophylactic measures to control enterotoxaemia disease, is also known as 'pulpy kidney disease', caused by *Clostridium perfringens* type D infections in sheep and goats.

VACCINATION SCHEDULE

Primary Vaccination: 4 months of age and above
Booster Dose: 6 months after primary vaccination
Re-Vaccination: Annual

PRECAUTIONS

Immunize only healthy animals. Malnutrition, infestation with worms, administration of immunosuppressive agents such as corticosteroids or radiotherapy will interfere with the immune response to vaccine.

POST-VACCINATION REACTIONS

Generally, there are no adverse reactions noticed. Occasionally a transient palpable swelling may occur at the site of injection, which will subside in 24-48 hours.

STORAGE & TRANSPORT

Store and transport the vaccine between 2 and 8°C until use. **At no stage should the vaccine be allowed to freeze.** Protect from sunlight.

DOSAGE & ROUTE

Sheep & Goats: 2 mL;
The vaccine should be injected subcutaneously in the mid neck region.

PRESENTATION

Pack: 100 mL



CORPORATE OFFICE: 6-2-1012, TGV Mansion, 5th Floor, Khairatabad, Hyderabad - 500 004
Telangana, India T: +91 40 6666 7464 / 6677 2726 F: +91 40 6610 4915 / 6677 2725

FACTORY: Plot Nos. 97, 98, 276 & 277, IDA, Pashamylaram, Patancheru, Sangareddy - 502 307, Telangana, India
T: +91 84552 24200 / 84552 24202 F: +91 84552 24201