

HAEMORRHAGIC SEPTICAEMIA VACCINE IP

HS VAC[®]

एचएस वैक

COMPOSITION

Each dose of HS VAC *, Haemorrhagic Septicaemia Vaccine contains formalin killed cultures of *Pasteurella multocida* (strain P52) adjuvanted with aluminium hydroxide gel. Formaldehyde (≤0.05% w/v) is added as a preservative.

INDICATIONS

HS VAC* is used for prophylactic measures to control haemorrhagic septicaemia disease caused by *Pasteurella multocida* infections in cattle and buffaloes.

VACCINATION SCHEDULE

Primary Vaccination: 6 months of age and above Booster Dose: 6 months after primary vaccination Re-Vaccination: Annual

PRECAUTIONS

Immunize only healthy animals. Malnutrition, infestation with worms, administration of immunosuppressive agents such as corticosteroids or radiotherapy will interfere with the immune response to vaccine.

POST-VACCINATION REACTIONS

Generally, there are no adverse reactions noticed. Occasionally a transient palpable swelling may occur at the site of injection, which will subside in 24-48 hours.

STORAGE & TRANSPORT

Store and transport the vaccine between 2 and 8°C until use. At no stage should the vaccine be allowed to freeze. Protect from sunlight.

DOSAGE & ROUTE

Cattle and Buffaloes: 2 mL;

The vaccine should be Injected subcutaneously in the mid neck region.

PRESENTATION

Pack: 100 mL





CORPORATE OFFICE: 6-2-1012, TGV Mansion, 5th Floor, Khairatabad, Hyderabad - 500 004 Telangana, India T: +91 40 6666 7464 / 6677 2726 F: +91 40 6610 4915 / 6677 2725

FACTORY: Plot Nos. 97, 98, 276 & 277, IDA, Pashamylaram, Patancheru, SangaReddy - 502 307 Telangana, India T: +91 84552 24200 / 84552 24202 F: +91 84552 24201