



RABIES VETERINARY VACCINE (CELL CULTURE), INACTIVATED IP

Rabivac Vet[®]

रेबीवैक वेट

COMPOSITION

Rabivac Vet[®] is used for prophylactic and post-bite treatment. It contains cell culture adapted rabies virus (Strain PV11) propagated on BHK-21 cell line and inactivated with aziridine compound. Aluminium hydroxide gel is used as an adjuvant. Each dose of 1 mL of Rabivac Vet contains potency \geq either 1.0 or 2.5 IU meeting the WHO standards.

INDICATION

The vaccine is intended for immunization of dogs and other domestic animals against rabies for prophylaxis and post-bite therapy.

PROPHYLACTIC USE

A single dose for animals of 3 months of age and above. Booster dose in 21 days after primary vaccination. Annual booster dose is recommended in endemic areas.



DOSAGE & ROUTE

Subcutaneous or intramuscular injection
Prophylactic. Dogs and all other species - 1 mL

POST-BITE THERAPY

1st Dose : On "0" day
(1st injection after the bite, preferably the day of bite)
2nd Dose : 3rd day
3rd Dose : 7th day
4th Dose : 14th day
5th Dose : 28th day

PRECAUTIONS

Immunize only healthy animals. Malnutrition, infestation with worms, administration of immunosuppressive agents such as corticosteroids or radiotherapy will interfere with the immune response.

STORAGE & TRANSPORTATION

Store and transport the vaccine at 2°C to 8°C. The antigenicity of the vaccine deteriorates if the temperature is allowed to rise above this range. At no stage should the vaccine be allowed to freeze.

PRESENTATION

Pack: 1 mL | 5 mL | 10 mL

CORPORATE OFFICE: 6-2-1012, TGV Mansion, 5th Floor, Khairatabad, Hyderabad - 500 004
Telangana, India T: +91 40 6666 7464 / 6677 2726 F: +91 40 6610 4915 / 6677 2725

FACTORY: Plot Nos. 97, 98, 276 & 277, IDA, Pashamylaram, Patancheru, Sangareddy - 502 307, Telangana, India
T: +91 84552 24200 / 84552 24202 F: +91 84552 24201