BLACKQUARTER
VACCINE IP

BQ VAC®

बीक्यू वैक

Brilliant
BIO PHARMA

Composition
Each dose of BQ VAC® Blackquarter Vaccine contains formalin killed cultures of *Clostridium chauvoei* (strain 49) adjuvanted with aluminium hydroxide gel. Formaldehyde (≤0.05% w/v) is added as a preservative.

Indications
BQ VAC® is used for prophylactic measures to control blackquarter disease caused by *Clostridium chauvoei* infections in cattle and buffaloes.

Vaccination Schedule
Primary Vaccination: 6 months of age and above
Booster Dose: 6 months after primary vaccination
Re-Vaccination: Annual

Precautions
Immunize only healthy animals. Malnutrition, infestation with worms, administration of immunosuppressive agents such as corticosteroids or radiotherapy will interfere with the immune response to vaccine.

Post-vaccination Reactions
Generally, there are no adverse reactions noticed. Occasionally a transient palpable swelling may occur at the site of injection, which will subside in 24-48 hours.

Storage & Transportation
Store and transport the vaccine between 2°C to 8°C until use. At no stage should the vaccine be allowed to freeze. Protect from sunlight.

Dosage & Administration
Cattle and Buffaloes: 2 mL: The vaccine should be injected subcutaneously in the mid-neck region.

SHAKE WELL BEFORE USE



PRESENTATION
Pack: 100 mL

