HAEMORRHAGIC
SEPTICAEMIA VACCINE IP

HS VAC®

एचएस वैक

Brilliant BIO PHARMA

Composition
Each dose of HS VAC®, Haemorrhagic Septicaemia Vaccine contains formalin killed cultures of *Pasteurella multocida* (strain P52) adjuvanted with aluminium hydroxide gel. Formaldehyde (≤0.05% w/v) is added as a preservative.

Indications
HS VAC® is used for prophylactic measures to control haemorrhagic septicaemia disease caused by *Pasteurella multocida* infections in cattle and buffaloes.

Vaccination Schedule
Primary Vaccination: 6 months of age and above;
Booster Dose: 6 months after primary vaccination
Re-Vaccination: Annual

Precautions
Immunize only healthy animals. Malnutrition, infestation with worms, administration of immunosuppressive agents such as corticosteroids or radiotherapy will interfere with the immune response to vaccine.

Post-Vaccination Reactions
Generally, no adverse reactions are noticed. Occasionally, a transient palpable swelling may occur at the site of the injection, which will subside in 24-48 hours.

Storage & Transportation
Store and transport the vaccine between 2°C to 8°C until use. At no stage should the vaccine be allowed to freeze. Protect from sunlight.

Dosage & Administration
Cattle and Buffaloes: 2 mL
The vaccine should be injected sub-cutaneously in the midneck region.

SHAKE WELL BEFORE USE



PRESENTATION
Pack: 100 mL

